

Body image dissatisfaction in patients with inflammatory bowel disease

Beese, Sophie; Harris, Isobel; Dretzke, Janine; Moore, David

DOI:
[10.1136/bmjgast-2018-000255](https://doi.org/10.1136/bmjgast-2018-000255)

License:
None: All rights reserved

Document Version
Peer reviewed version

Citation for published version (Harvard):
Beese, S, Harris, I, Dretzke, J & Moore, D 2019, 'Body image dissatisfaction in patients with inflammatory bowel disease: a systematic review', *BMJ Open Gastroenterology*, vol. 6, no. 1, e000255.
<https://doi.org/10.1136/bmjgast-2018-000255>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:
Checked for eligibility 06/02/2019

This is an author-produced, peer-reviewed version of an article forthcoming in *BMJ Open Gastroenterology*.
<https://bmjopengastro.bmj.com/>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BMJ Open Gastroenterology

Body Image Dissatisfaction in Patients with Inflammatory Bowel Disease: A Systematic Review

Journal:	<i>BMJ Open Gastroenterology</i>
Manuscript ID	bmjgast-2018-000255.R1
Article Type:	Systematic review
Date Submitted by the Author:	19-Nov-2018
Complete List of Authors:	Beese, Sophie; University of Birmingham, Institute of Applied Health Research Harris, Isobel; University of Birmingham, Institute of Applied Health Research Dretzke, Janine; University of Birmingham, Institute of Applied Health Research Moore, David; University of Birmingham, Institute of Applied Health Research
Keywords:	INFLAMMATORY BOWEL DISEASE, QUALITY OF LIFE, ULCERATIVE COLITIS, CROHN'S DISEASE
Abstract:	<p>Background and Aims: Little is known about the relationship between inflammatory bowel disease and body image. The aim of this systematic review was to summarise the evidence on body image dissatisfaction in inflammatory bowel disease patients across four areas: body image tools (i), prevalence (ii), factors associated with body image dissatisfaction in inflammatory bowel disease (iii) and association between inflammatory bowel disease and quality of life (iv).</p> <p>Methods: Two reviewers screened, selected, quality assessed and extracted data from studies in duplicate. EMBASE, MEDLINE, PsycINFO and Cochrane CENTRAL were searched to April 2018. Study design specific critical appraisal tools were used to assess risk of bias. Narrative analysis was undertaken due to heterogeneity.</p> <p>Results: Fifty-seven studies using a body image tool were included; 31 for prevalence and 16 and 8 for associated factors and association with quality of life respectively. Studies reported mainly mean or median scores. Evidence suggested female gender, age, fatigue, disease activity, and steroid use were associated with increased body image dissatisfaction, which was also associated with decreased quality of life.</p> <p>Conclusion: This is the first systematic review on body image in inflammatory bowel disease patients. The evidence suggests that body image dissatisfaction can negatively impact patients, and certain factors are associated with increased body image dissatisfaction. Greater body image dissatisfaction was also associated with poorer quality of life.</p> <p>However, the methodological and reporting quality of studies was in some cases poor with considerable heterogeneity. Future IBD research should incorporate measurement of body image dissatisfaction using validated tools.</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

**Body Image Dissatisfaction in Patients with Inflammatory Bowel Disease:
A Systematic Review**

**Sophie Elizabeth Beese¹, MPH, Isobel Marion Harris¹, MPH, Janine
Dretzke¹, MSc, David Moore¹, PhD.**

*¹Institute of Applied Health Research, Public Health Building, University of
Birmingham, B15 2TT. 0121 415 9107.*

Corresponding author: Sophie Beese, s.beese@bham.ac.uk

Word Count: 3694

ABSTRACT

Background and Aims: Little is known about the relationship between inflammatory bowel disease and body image. The aim of this systematic review was to summarise the evidence on body image dissatisfaction in inflammatory bowel disease patients across four areas: body image tools (i), prevalence (ii), factors associated with body image dissatisfaction in inflammatory bowel disease (iii) and association between inflammatory bowel disease and quality of life (iv).

Methods: Two reviewers screened, selected, quality assessed and extracted data from studies in duplicate. EMBASE, MEDLINE, PsycINFO and Cochrane CENTRAL were searched to April 2018. Study design specific critical appraisal tools were used to assess risk of bias. Narrative analysis was undertaken due to heterogeneity.

Results: Fifty-seven studies using a body image tool were included; 31 for prevalence and 16 and 8 for associated factors and association with quality of life respectively. Studies reported mainly mean or median scores. Evidence suggested female gender, age, fatigue, disease activity, and steroid use were associated with increased body image dissatisfaction, which was also associated with decreased quality of life.

Conclusion: This is the first systematic review on body image in inflammatory bowel disease patients. The evidence suggests that body image dissatisfaction can negatively impact patients, and certain factors are associated with increased body image dissatisfaction. Greater body image dissatisfaction was

also associated with poorer quality of life. However, the methodological and reporting quality of studies was in some cases poor with considerable heterogeneity. Future IBD research should incorporate measurement of body image dissatisfaction using validated tools.

Keywords: Systematic review – inflammatory bowel disease – body image – quality of life

What is already known about this subject?

Body image in inflammatory bowel disease (IBD) patients is relatively unexplored. However, it may potentially be an issue as symptoms and other factors associated with IBD could lead to body image dissatisfaction. Both of these conditions may lead to an increased risk of developing mental health disorders such as depression and self-harm, warranting research into the area.

What are the new findings?

This review highlights associations between certain factors in IBD and body image dissatisfaction, as well as suggesting that increased body image dissatisfaction is related to reduced quality of life. The review also highlights the need for a validated, reliable tool to measure body image in IBD patients as well as clearer reporting of studies.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

How might it impact on clinical practice in the foreseeable future?

Body image can be considered when making clinical decisions regarding IBD patients with the potential for interventions to improve body image in patients at risk.

81 INTRODUCTION

82

83 Inflammatory bowel disease (IBD) is associated with a range of debilitating
84 symptoms⁽¹⁾ and affects around 300,000 people in the UK⁽²⁾, over 1 million in
85 the USA and 2.5 million across Europe⁽³⁾. A potentially overlooked issue for IBD
86 patients is body image dissatisfaction (BID). Body image (BI) is how an
87 individual perceives themselves physically⁽⁴⁾ and sufferers have a distorted and
88 negative view of themselves, feeling anxious and uncomfortable about their
89 body. Additionally, negative BI can have a serious impact on health and well-
90 being⁽⁵⁾.

91 Social media and celebrity attention contribute to pressure to adhere to an
92 “ideal” body and an obsession with appearance^(6, 7). Discontentment with
93 aspects such as body weight, shape, appearance and skin may contribute
94 toward an individual having BID⁽⁸⁾. Studies have shown patients with negative
95 BI are more likely to suffer with depression, anxiety and feel suicidal and BID
96 can impact negatively upon relationships⁽⁹⁾ and quality of life (QoL)⁽¹⁰⁾.

97 Various tools have been utilised in healthcare to measure BI including the Body
98 Image Ideals Questionnaire, the Body Image Scale and the Cash Body Image
99 Disturbance Questionnaire (BIDQ)⁽¹¹⁾. There are also condition-specific BI tools
100 such as the Body Image Scale (BIS) for IBD⁽¹²⁾.

101 Both condition-specific symptoms and treatments may contribute to BID in IBD
102 patients, particularly during periods of active disease rather than remission.

103 Symptoms can include urgent bowel movements, bloating, excess wind, fatigue,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

104 skin problems and ulcers. Treatment with steroids can be associated with
105 weight gain, acne and mood swings⁽¹³⁾. Surgeries may also impact upon BI due
106 to scarring and implementation of a stoma⁽¹⁴⁾ ⁽¹⁵⁾. Those suffering with IBD or
107 BID are at an increased risk of mental health issues^(16, 17) ; this could be worse
108 for patients living with both conditions. Furthermore, most IBD patients are
109 diagnosed at adolescence⁽¹⁸⁾, when BI is important. Body image is currently not
110 routinely considered in the management of IBD.

111 No existing or ongoing systematic reviews on BI in IBD have been identified.
112 However multiple primary studies, mainly cross-sectional in nature assess BI as
113 an outcome in IBD patients, with disparate results. A systematic review is
114 therefore warranted to synthesise and clarify the evidence base.

115
116 The following four questions will be addressed:

- 117
- 118 **1.** What tools are used to measure body image in IBD patients and what are
 - 119 their components?
 - 120 **2.** What is the prevalence and severity of body image dissatisfaction in IBD
 - 121 patients?
 - 122 **3.** What factors are associated with body image dissatisfaction in IBD patients?
 - 123 **4.** Is there an association between body image dissatisfaction in IBD patients
 - 124 and quality of life?

METHODS

This systematic review has been reported according to the Preferred Reporting Items of Systematic Reviews and Meta-Analysis (PRISMA) guidelines⁽¹⁹⁾. A protocol was previously registered (PROSPERO (CRD42018060999)) and submitted for publication and is currently in process⁽²⁰⁾. A summary of the methods is reported below. Selection, data extraction and quality assessment were carried out by two independent reviewers with disagreements resolved through discussion or third reviewer.

Search Strategy

Bibliographic databases (EMBASE, MEDLINE, PsycINFO, Cochrane CENTRAL) were searched to April 2018 using combinations of index and text terms for IBD and BI (see Supplementary table 1 for MEDLINE strategy). Strategies were adapted for each database and run without date or language restrictions. Trial registries (Clinicaltrials.gov, EU Clinical Trial Register) were searched for ongoing trials and reference lists of included studies were checked.

Screening and Selection Criteria

Study eligibility was based on the following criteria:

Study design: Any primary study reporting quantitative data.

Population: Patients of any age diagnosed with IBD. At least 50% of population must have IBD unless results are reported separately for sub-groups of individuals with IBD.

Tools: Any tool measuring any aspect of BI (including QoL tools that had at least one BI related domain or question).

Studies were also eligible (for question 2-4) where they reported: any measure of prevalence/frequency and severity of BID in IBD patients; data on associations between any factor in IBD patients and BID; or any association between BI and QoL measures in IBD patients, including associations between two separate domain measures of the same tool.

Exclusion criteria: Case reports, qualitative research and conference abstracts published three years before the date of the searches.

Reasons for exclusion were recorded.

159

Data Extraction

A piloted data extraction form was used. Examples of the type of data extracted are shown below:

Study characteristics:

Study design, aim and setting, inclusion/exclusion criteria, recruitment methods, follow-up period.

Participant characteristics:

167 Number of patients, age, gender, type of IBD, disease severity and activity,
168 BMI, comorbidities, therapy/surgery.

169 Data for synthesis/analysis:

170 BI measurement tool, components of tools/scales, data on BID (e.g. body image
171 scores, prevalence, thresholds for determining BID), factors associated with BI
172 dissatisfaction and strength of association, QoL measures, strength of
173 association between BID and QoL.

174

175 **Quality Assessment**

176 Quality assessment was based on critical appraisal checklists for both
177 prevalence and cross-sectional analytical studies from the Joanna Briggs
178 Institute⁽²¹⁾. Studies solely included for question one were not quality assessed
179 as the objective of this question was to compile a list of body image tools.

180 Important quality items included sample selection, response rate during
181 enrolment in the study, clear inclusion criteria and measurement of outcomes in
182 a valid and reliable way.

183

184 **Analysis**

185 A narrative synthesis was carried out separately for each question, with key
186 findings tabulated. Substantial heterogeneity relating to populations, tools and
187 settings was apparent in the included studies meaning that meta-analysis was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

188 not appropriate. Consistencies and discrepancies in findings between studies
189 were noted and discussed in the context of any likely sources of heterogeneity.
190 Quality assessment findings were used when considering the strength of
191 evidence for the latter three questions.

Confidential: For Review Only

RESULTS

Database searches identified 587 records and 57^(14, 22-77) studies were included, with some studies eligible for multiple questions (see Figure 1 for selection process and reasons for exclusion). All 57 papers reported using BI tools, 31^(14, 22-26, 30, 31, 33-39, 42, 47, 50, 51, 53, 54, 59, 61, 62, 64-66, 68, 70, 72, 73) reported prevalence or mean/median BI scores, 16^(14, 23, 24, 30, 34-36, 47, 54, 59, 61, 62, 64, 66, 68, 72) studies presented factors associated with BID and 8^(14, 22-24, 34, 62, 66, 72) studies reported correlations between QoL and body image.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

227 **Question 1: What tools are used to measure body image and what are**
228 **their components?**

229 Of the 57 studies measuring BI, 51 were cross-sectional whilst the others varied
230 (case-control⁽²⁵⁾, prospective cohort^(51, 66), case series⁽³⁹⁾, randomised controlled
231 trial⁽⁶⁵⁾ and non-randomised intervention study⁽⁴²⁾). Study populations included
232 adults and children in settings including outpatients, pre/post-surgery, summer
233 camps and online registries, from countries across the world. Twenty studies
234 focused on BI as one of the main outcomes but only six of these studies were
235 non-surgery based.

236 Fifteen tools were identified (Table 1). Seven tools were specifically for BI and
237 eight were QoL tools which included a BI domain or question(s). The most
238 frequently applied tool specific to BI was the Body Image Questionnaire (BIQ)
239 which was used in 14 studies. The Body Image Scale (BIS) was used in 5
240 studies and is the only tool validated in an IBD population. IMPACT-III (or earlier
241 IMPACT-II) is a validated QoL questionnaire aimed at adolescents and children
242 with IBD and includes a BI domain. It was used across 18 studies. The
243 remaining 12 tools were used in only 1-3 studies respectively.

244 None of the tools included had a clear cut-off point for defining BID but offered
245 an indication of increasing or decreasing likelihood of dissatisfaction. In some
246 tools, a higher score indicated better BI (BIQ, EORTC, DUX-25). In others, a
247 higher score indicated increased BID (IMPACT, BIS, RFIPC, IBDSI, Body
248 Image Self-Consciousness during Intimacy Scale, BIDQ and ASWAP).

249 Tools where items had similar themes were grouped to show general focus of
 250 BI questions and are shown in Table 2.

251

Table 1. Tools identified and used across included studies

Measurement Tool	Type of Tool	Intended target population	Is tool Validated?	Scoring	No. of Studies Tool Used In
Body Image Tools					
ASWAP	Body image	Initially used in scleroderma patients	Yes but not in IBD patients	15 items rated on 7-point scale. Questions corresponding to items 4–11 were reverse scored such that higher scores reflect greater dissatisfaction	1
Askevold's Body Image Test	Body image	Unclear	Unclear	Unclear	2
Body Image and Self-Consciousness During Intimacy Scale	Body image and sexual self-consciousness	Women	No	0-75, higher scores poorer body image	1
BIA/BIA-P	Body image	Adults, no specific clinical population	Unclear	Based on body image silhouettes ranging in size. Score=difference between current body size and ideal body size.	1
BIQ	Body image	Originally caesarean or appendectomy patients, now IBD	No	5-20, higher score better body image	14

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

		patients			
BIS	Body image	Cancer patients	Yes	0-30, lower score better	5
				body image	
Cash Body Image Disturbance Questionnaire	Body image	Range of clinical groups	Yes but not in IBD patients	7-35, higher score poorer body image	2
Quality of Life Tools with a Body Image Component					
BUX-25	Quality of daily functioning. (1 of 4 domains relate to body image)	School age children	No	Higher scores, better QoL	1
EORTC-QLQ-CR38	Quality of Life questionnaire. (3 of 38 items relate to body image)	Cancer patients	Yes but not in IBD patients	38 items with 4 category responses. Functional scales: higher score higher functioning. Symptoms scales: higher score higher level of symptoms.	1
EORTC-QLQ-CR29	Quality of Life questionnaire. (3 of 29 items relate to body image)	Cancer patients	Yes but not in IBD patients	29 items with 4 category responses. Functional scales: higher score higher functioning. Symptoms scales: higher score higher level of symptoms.	1
IMPACT-III or IMPACT II	Health-related quality of life. (3 of 35 items relate to body image)	Children and adolescents with IBD	Yes	35-175, higher scores better QoL	18
Inflammatory Bowel Disease Stress Index	Assessing the extent to which IBD has caused alterations in lifestyle. (1 of 10 items relate to body image)	IBD patients	Unclear	8 scales with a score of 0-3 (no impact-a great deal of impact).	1

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

image)

RFIPC	Quality of life questionnaire. (1 item of 25 relate to body image)	IBD patients	Yes	0-100, higher score poorer QoL	3
Stoma Quality of Life Scale	Stoma-related. (5 items of 19 relate to body image and sexuality).	Stoma patients	Yes (in ostomy patients)	5 scales, 19 questions. Each scored 1-5 (Never-always). Average scores for each scale calculated.	3
The Karolinska Psychodynamic Profile	Assessment of stable modes of mental functioning and character traits (1 subscale and 3 of 18 items relate to body image.)	No specific clinical population	Yes	Each subscale is graded from 1-3 (Most normal-least normal).	2

Abbreviations: ASWAP: Adapted Satisfaction with Appearance scale; BI/BIA-P: Body Image Assessment/Body Image Assessment-Preadolescent; BIQ: Body Image Questionnaire; BIS: Body Image Scale; DUX-25: Dutch Children's AZL/TNO Quality of Life Questionnaire; EORT-QLQ-CR38/EORT-QLQ-CR29: European Organisation for Research and Treatment of Cancer (EORTC) Quality of Life questionnaire for Colorectal Cancer; IMPACT-II/IMPACT-III: A measure of health-related quality of life in paediatric inflammatory bowel disease; RFIPC: Rating Form of IBD Patient Concerns.

252

253

254

255

256

257

258

Table 2. Body image tools with similar questions grouped into overarching themes

Body Image Tool	Components								
	Satisfaction with appearance	Attractiveness	Socialising/Work	Avoidance of people or tasks	Feeling feminine/masculine	Effect of disease on body	Scar satisfaction	Satisfaction with body both naked and dressed	Distressing thoughts
BIS	✓	✓	✓		✓	✓		✓	
BIQ	✓	✓			✓	✓	✓	✓	
CBIDQ	✓	✓	✓	✓					✓
ASWAP	✓	✓	✓	✓				✓	
N.B. Similar components of tools were grouped into themes shown above. BIS=Body Image Scale. BIQ=Body Image Questionnaire.									

CBDIQ=Cash Body Image Disturbance Questionnaire. ASWAP=Adapted Satisfaction with Appearance Scale. Askevold's Body Image Test (no information in paper or online), Body Image and Self-consciousness during Intimacy Scale (too specific) and the Body Image Assessment (based on figural drawing scales) were not included.

What is the prevalence of body image dissatisfaction in IBD patients?

Thirty-one studies including a total of 3,634 patients reported on prevalence or severity of BID (see Table 3 for study characteristics). Seventeen studies^(14, 22, 23, 25, 30, 31, 38, 42, 53, 54, 59, 61, 62, 66, 70, 72, 73) included both ulcerative colitis (UC) and Crohn's disease (CD) patients. Ages ranged from 2-71 and eighteen studies^(22, 30, 38, 40-42, 51-53, 58, 60-63, 70, 71, 73, 76) included only children/adolescents. Fourteen studies^(24-26, 33-37, 39, 47, 50, 64, 65, 68) included surgery patients and one study included only females⁽⁷²⁾.

Only three studies reported prevalence. Brown (2015)⁽²⁶⁾ found that 21-34% UC patients reported negative impacts on BI using BIQ. McDermott (2015)⁽¹⁴⁾ found that 87% patients reported some form of concern about an aspect of their BI using the Cash Body Image Disturbance Questionnaire. Muller (2010)⁽⁵⁹⁾ reported that 66.8% IBD patients stated they had impaired BI based on a researcher devised questionnaire. The other 28 studies reported mean/median BI scores based on a range of tools.

In studies with populations undergoing surgery it was found that there was no significant difference in BI scores (using the BIQ) after laparoscopic or open/conventional surgery in IBD patients^(33-35, 64, 78). Only one study found BI scores to be significantly improved after laparoscopic surgery compared to conventional surgery in CD⁽³⁶⁾.

BI was included as an outcome across 31 studies. All but one study compared results within the included IBD population e.g. UC vs CD, surgery vs no surgery, males vs females. Bel (2015) found that women with IBD with disease in

remission scored comparably to women in a healthy population. One longitudinal study by Saha (2015)⁽⁶⁶⁾ measured scores over two years and found that BI did not change despite improvements in symptoms.

Confidential: For Review Only

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Beld et al (2010)	Cross sectional	UC or FAP undergone restorative proctocolectomy IPAA Jan 92 to Oct 08	Netherlands	26	UC (16) FAP (10)	BIQ	Mean body image scores (SD)	Males 16.3 (3.1) Females 13.5 (4.1)
Brown et al (2015)	Cross sectional	Patients with UC who had colectomy within the past 10 years, data collected from Nov 2010 to July 2011.	Canada, Australia, UK	351	All UC	BIQ	Median body image scores (IQR) Prevalence of “quite a bit” or “extreme” negative impacts on body image as a result of colectomy.	Males 8 (IQR 6-11) Females 11 (IQR 8-14) Age group >50 years 8 (IQR 6-11) Age group <50 years 10 (IQR 7-13). 21-34% reported negative impacts on body image.
Dunker et al (1998)	Cross sectional	CD patients undergoing open or laparoscopic resection at Leiden university medical centre	Netherlands	34	All CD	BIQ	Mean body image scores	Open 16.4 (10-20) Laparoscopic 18 (13-20) (SD not reported)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Dunker et al (2001)	Cross sectional matched comparison	UC patients who underwent laparoscopic assisted IPAA and matched conventional IPAA patients.	Netherlands	32	UC (28) FAP (4)	BIQ	Mean body image scores (SD)	Laparoscopic 19 (1.3) Conventional 17.9 (SD not reported)
Eshuis et al (2008)	Repeated cross sectional	Patients who underwent ileocolic resection for Crohn's disease from 1995 until 1998 two centres.	Netherlands	71 (medical file analysis) 61 (returned questionnaires)	All CD	BIQ	Mean body image scores (range)	Open 15.63 (6-20) Laparoscopic 16.3 (7-20) (SD not reported)
Eshuis et al (2010)	Repeated cross sectional	CD patients who had ileocolic resections between September 1999 and November 2003.	Netherlands	55	All CD	BIQ	Median body image scores (IQR)	Open 18.0 (IQR 16-19) Laparoscopic 19.0 (IQR 17-20)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Giudici et al (2017)	Case series (Abstract only)	December 2014-December 2015. Consecutive patients undergoing laparoscopic proctectomy for ulcerative colitis.	Italy	10	All UC	Self-designed body image questionnaire	Mean body image score	59 (SD not reported)
Kjaer et al (2014)	Cross sectional	Adult patients treated with laparoscopy-assisted or open IPAA at Odense University Hospital during the period between October 2008 and March 2012.	Denmark	50	UC (44) FAP (4) Other (2)	BIQ	Median body image scores (range)	Laparoscopic 8 (5-18) Open 9.5 (5-20)
Polle et al (2007)	Repeated cross sectional	Patients eligible for an elective proctocolectomy with IPAA for UC or FAP were included in a randomized trial	Netherlands	53	UC (34) FAP (19)	BIQ	Mean body image scores (limited data)	Women open group: 15 Laparoscopic group: 18 (SD not reported)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Ponsioen et al (2017)	Randomised controlled trial	Eligible patients aged 18–80 years, had active Crohn's disease of the terminal ileum, and had not responded to at least 3 months of conventional therapy with glucocorticosteroids, thiopurines, or methotrexate. Patients with diseased terminal ileum longer than 40 cm or abdominal abscesses were excluded.	Netherlands and UK	70 patients Infliximab group 73 Laparoscopic ileocaecoresection	All CD	BIQ	Mean body image scores (only given for resection group)	Resection group: Baseline 16. Endpoint 17.8. (SD not reported)
Scarpa et al (2009)	Prospective case series	Patients admitted for intestinal surgery for CD May 06 - July 08	Italy	47	All CD	BIQ	Median body image score (IQR)	5 (5-8)
Voermans et al (2010)	Prospective case series	A consecutive series of patients who had an indication for a laparoscopic ileocolic resection were invited to participate. CD patients.	Netherlands	10	All CD	BIQ	Median body image scores	Before surgery 17.0 After surgery 19.0
Bengtsson et al (2011)	Case-control	Patients with preoperative diagnosis of UC or CD who underwent IPAA	Sweden	101 (72 controls,	Controls; UC (60) CD (0) Study group;	BIS	Median body image scores.	Study group: Males 6.5 Females 10. Control group Males

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
				29 study group)	UC (25) CD (4)			1 Females 3
Trindade et al (2017)	Cross sectional	Female participants with ages between 18 and 40 years old who had not undergone IBD-related surgery	Portugal	96	UC (58) CD (38)	BIS	Mean body image score (SD)	10.10 (7.73) (SD not reported)
Vlahou et al (2008)	Cross sectional	Adolescents with IBD who attended clinics at two separate hospitals and a camp for children with IBD	USA	44	Breakdown not reported	BSQ (modified version of BIQ) and BIA-P	Mean body image scores (SD)	BSQ: Males 36.45 (4.88) Females 33.52 (7.77). BIA-P: Males 0.41 (0.85) Females 0.77 (0.92)
Grootenhuys (2009)	Non-randomised controlled	Adolescents with IBD who were under medical care at Emma Children's Hospital AMC and members of Crohn's and colitis association Netherlands	Netherlands	18 controls; 22 interventi	Controls CD (11) UC (4) IBDU (3). Intervention	DUX-25	Mean body image domain scores (SD)	Intervention: baseline 55.4 (18.6) post intervention 68.9 (17.7) Control:

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
	study			on	CD (17) UC (5) IBDU (0)			baseline 60.0 (17.4) post intervention 59.0 (20.1)
Bel et al (2015)	Cross sectional with controls	18-70 UC or CD	Netherlands	287 (197 healthy controls)	UC (132) CD (155)	EORTC-QLQ-CR38	Mean body image domain scores (SD)	Active: Males 5.61 (2.31) Females 6.2 (2.78). Remission: Males 3.82 (1.33) Females 4.58 (1.68)
Shepanksi (2009)	Before and after study	Children attending Camp Guts and Glory in Pennsylvania	USA	61	CD:UC (2:1)	IMPACT II	Mean body image domain scores (SD, for before and after camp)	By age; Age 9-10: pre 14.6 (4.1). Post 16.4 (3.7). Age 11-12: Pre 11.4 (4.9). Post 13.2 (5.0). Age 13-14: Pre 12.9 (5.2). Post 13.8 (5.9). Age 15-16: Pre 12.3

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
								(5.0). Post 11.2 (5.4)
Abdovic et al (2013)	Cross sectional validation study	Children aged nine years or older with confirmed diagnosis of IBD for more than six months from inpatient and outpatient clinics at particular centres.	Croatia	104	UC (30) CD (74)	IMPACT III	Mean body image domain score (SD).	12.03 (1.96)
Chouliara s et al (2017)	Cross sectional	UC and CD patients hospitalized or followed in outpatient clinic in Athens	Greece	99	UC (37) CD (62)	IMPACT III	Mean body image domain scores (SD)	Overall 71.5 (17.9) UC 67.3 (22.4) CD 72.6 (19.3) No significant relationship between body image and assessed disease characteristics or

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
								prescribed medications.
Gallo et al (2014)	Cross sectional	Children between the ages of 8 and 18 years, who had been diagnosed with IBD at least 6 months before, and were being followed at the Pediatric Gastroenterology Service of the Hospital Italiano de Buenos Aires, Argentina, or at the private office of one of the co-authors (M.O.) and one of their parents.	Argentina	27	UC (17) CD (9)	IMPACT III	Mean body image domain score (SD)	76.54 (16.06)
Lee et al (2015)	Prospective observational study	Children and young adults less than 22 years of age started on EN or anti-TNF therapy for active CD at Hospital for Sick Children Toronto and Children's Hospital Philadelphia.	Canada and USA	90	All CD	IMPACT III	Median body image domain scores (range)	Baseline PEN 71 (54-75) EEN 58 (58-75) TNf 67 (50-83)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Mason et al (2015)	Prospective observational study	Adolescents >10 years old with confirmed diagnosis of IBD attending gastroenterology clinic at Royal Hospital for Sick Children, Glasgow	Scotland	63	UC/IBDU (18) CD (45)	IMPACT III	Mean body image domain score	7 (SD not reported)
Ogden et al (2011)	Cross sectional validation study	Unclear - children with IBD	UK	97	UC (12) CD (64) IBDU (21)	IMPACT III	Mean body image domain score	63.5 (95% CI 56.5 - 70.6) (SD not reported)
Perrin et al (2008)	Cross sectional	Children aged 8-17 years diagnosed with UC or CD 6 mnths before the study followed at 1 of 6 paediatric gastroenterology centres. No other chronic conditions.	USA	220	UC (59) CD (161)	IMPACT III	Mean body image domain scores (SD)	68.1 (19.6) UC 68.6 (20.8) CD 67.9 (19.2)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
McDermot et al (2015)	Cross sectional	Patients with histologically confirmed IBD attending ambulatory clinics in 1 of 2 medical centres between July 2011 and November 2012	Ireland	330	UC (145) CD (194)	Modified BIS and Cash Body Image Scale (qualitative only)	Median body image score (range) Prevalence	6 (0-27) 13% patients reported no concerns about any aspect of body image
Saha et al (2015)	Prospective observational study	Patients with UC, CD or IBDU aged 18 and above enrolled in the Ocean State Crohn's and Colitis Area Registry (OSCCAR) with a minimum of 2 years of follow-up	USA	274	CD (145) UC/IBDU (129)	ASWAP	Mean body image scores (SD)	Baseline: Females 30.1 (14.4) Males 21.2 (8.4) Year 1: Females 28.2 (14.1) Males 24.5 (12.5) Year 2: Females 28.8 (13.2) Males 24.1 (13.5)

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
Muller et al (2010)	Cross sectional	IBD patients aged 18-50 from a database of patients maintained by the Southern Adelaide IBD Service	Australia	217	UC (85) CD (127) IBDU (5)	No specific tool – range of questions regarding body image and impact of IBD on this	Prevalence (%) of body image dissatisfaction	66.8% of patients reported impaired body image
de Rooy et al (2001)	Cross sectional	Outpatients of the Inflammatory Bowel Disease Center, Mount Sinai Hospital. Subjects were a convenience sample waiting for a regularly scheduled physician appointment.	USA	241	UC (121) CD (120)	RFIPC	“Feelings about body” question mean score (SD)	42.84 (33.97)
Maunder et al (1999)	Retrospective analysis	Patients with IBD who had completed the RFIPC and a survey of demographic and disease-related variables in one of three previous studies	Unclear	343	UC (186) CD (157)	RFIPC	“Feelings about body” question mean scores	Female 52.13 (34.8) Male 38.16 (33.83)
Kuruville et al	Cross-sectional	Consecutive patients who had undergone IPAA or a permanent ileostomy for ulcerative colitis by a	USA	59	All UC. IPAA (35); TPC (24).	Stoma Quality of Life Scale	Mean (SD) and median (range) body	IPAA; Mean 93.1 (9.7). Median 100.

Table 3. Study characteristics of papers included for questions two, three and four.

Study	Design	Population	Country	No. patients	No. UC/CD/Other	Body Image Tool	Outcomes	Body Image Prevalence/Score
(2012)	(Abstract only)	single surgeon, presenting for their annual follow-up visit from July through September 2011, were offered participation in the study. A randomly chosen group of subjects who did not have scheduled appointments during the study period were sent a letter inviting them to participate in the study.					image/sexuality domain scores.	(65-100). TPC: Mean 76.4 (14.6) Median 80 (50-100).
<p>Abbreviations: UC: Ulcerative Colitis; CD: Crohn's disease; IBD: Inflammatory bowel disease; IBDU: Inflammatory bowel disease unclassified; FAP: Familial adenomatous polyposis; IPAA: ileal pouch-anal anastomosis; TPC: Total proctocolectomy; PEN: Partial Enteral Nutrition; EEN: Exclusive Enteral Nutrition; TNF: Tumour Necrosis Factor; TNF: Tumour necrosis factor; IQR: Interquartile range; SD: Standard deviation; ASWAP: Adapted Satisfaction with Appearance scale; BI/BIA-P: Body Image Assessment/Body Image Assessment-Preadolescent; BIQ: Body Image Questionnaire; BIS: Body Image Scale; DUX-25: Dutch Children's AZL/TNO Quality of Life Questionnaire; EORT-QLQ-CR38/EORT-QLQ-CR29: European Organisation for Research and Treatment of Cancer (EORTC) Quality of Life questionnaire for Colorectal Cancer; IMPACT-II/IMPACT-III: A measure of health-related quality of life in paediatric inflammatory bowel disease; RFIPC: Rating Form of IBD Patient Concerns; BSQ: Body Satisfaction Questionnaire.</p>								

302

303

304

What factors are associated with body image dissatisfaction in IBD patients?

Sixteen studies^(14, 23, 24, 30, 34-36, 47, 54, 59, 61, 62, 64, 66, 68, 72) totalling 2333 IBD patients reported the association between various factors and BID (see Table 4).

Factors included those related to demographics as well as disease and treatment-related characteristics. Ten studies^(14, 24, 34-36, 47, 64, 66, 68, 72) utilised a specific BI tool and six^(34-36, 47, 64, 68) focused on comparative surgery techniques. Three studies^(30, 61, 62) included a paediatric population; the remaining studies included adults. BI was one of the main outcomes in most of these studies and the study by Saha (2015)⁽⁶⁶⁾ was the first longitudinal follow up of BID in IBD according to the authors.

In 6/10 studies^(14, 23, 54, 59, 64, 66) female gender was found to be significantly associated with increased BID. One study⁽⁵⁹⁾ reported the odds of BID was over 3 times more in females than males ($p=0.001$), with strong associations reported in the other five studies. Increased disease activity was found to have a significant but moderate positive association in 7/9 studies^(14, 23, 34, 62, 66, 68, 72)

Other factors found to be significantly associated with increased BID included steroid use^(14, 61, 66, 72), age^(14, 23), increased BMI^(14, 72), smoking⁽¹⁴⁾ and fatigue⁽²³⁾(Table 4). Saha (2015)⁽⁶⁶⁾ also found a significant association between extraintestinal manifestations (EIM) and increased BID, but were the only study to assess this. Laparoscopic surgery was found to be associated with improved body image in 2/6 studies^(36, 68). Ileal pouch-anal anastomosis (IPAA) seemed to result in patients being satisfied with their body image in two studies^(24, 26) but

they lacked a comparative surgery group. One study⁽⁵⁰⁾ compared IPAA and ileostomy and found better body image scores in the IPAA group. No significant associations were found between disease sub-type and increased BID.

Is there an association between body image dissatisfaction and quality of life in IBD patients?

Eight studies^(14, 22-24, 34, 62, 66, 72) explored a potential association between BID and QoL across a total of 1371 patients, with seven presenting a significant association. Three studies^(22, 24, 62) (Table 4) focused on younger populations with the rest including adults only. The majority of studies included populations with both UC and CD whilst two^(24, 34) included only one subtype.

Statistically significant weak to moderately strong correlations were present in five studies^(22, 23, 34, 62, 72) ranging from $r = 0.34$ to $r = 0.67$. Furthermore, McDermott(2015)⁽¹⁴⁾ found that when using the BI scale there was a significant difference in scores between those with good or poor QoL. Trindade(2017)⁽⁷²⁾ found that BI was positively correlated with psychological and physical QoL. Saha(2015)⁽⁶⁶⁾ found that a one unit increase in the total ASWAP score (indicating poorer body image) was associated with a 0.62 decrease in QoL score ($p < 0.0001$).

Various QoL tools (See Table 1) were used across studies with some using more than one. Four of these questionnaires used (IMPACT II and III, GIQLI and WHOQOL-BREF) contain a question or domain on BI, potentially making them more likely to correlate with BI questionnaires.

1
2
3
4 351
5
6
7 352
8
9 353
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Confidential: For Review Only

Table 4. Most common factors found to be significantly associated with impaired body image in IBD as reported in each study, including associations between reduced body image and reduced QoL.

	Abdovic	Bel 2015	Bel 2010	Chouliaras 2017	Dunker 1998	Eshuis 2008	Eshuis 2010	Kjaer 2014	Mander 1999	McDermott 2015	Muller 2010	Ogden 2004	Perrin 2007	Saha 2015	Scarpa 2009	Trindade 2017
Female gender	r= -0.18*	Difference in means p=0.08	Difference in means p>0.10	Difference in scores p=0.18	No significant association			Female	p<0.001*	Difference in proportions p=0.007		Significantly worse scores in open surgery group p=0.004*		p<0.0001*		

Factor	Study																
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	Dunker 1998	Eshuis 2008	Eshuis 2010	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Ogden 2000	Perrin 2000	Polle 2007	Saha 2015	Scarpa 2009	Trindade 2017
Higher disease/symptom activity	r=			No	r=0.5*					p<0.0		p=	p=		In UC	Multiple	Active
	0.3			signific						01*		0.5	0.00		p=0.006	regressi	diseas
	8*			ant								0	3*		*	on	e
				associ											In CD	β=0.426	r=0.18
				ation											p=0.003	p=0.006	
															*	*	Sympt
																	oms
																	r=0.40
																	*
Fatigue	r=																
	0.5																
	5*																

Factor	Study																
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	Dunker 1998	Eshuis 2008	Eshuis 2010	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Ogden 2000	Perrin 2000	Polle 2007	Saha 2015	Scarpa 2009	Trindade 2017
Disease Subtype				No significant association						p=0.63	Difference in proportions p=0.094		p=0.05		No association found		
Age		r= -0.18*			No significant association					Younger age p<0.001*							r= -0.06
Steroids				No significant association	No significant association					p=0.03*		p=0.005*			p=0.02*		r=0.22*

Factor	Study														
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	Dunker 1998	Eshuis 2008	Eshuis 2010	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Ogden 2000	Perrin 2000	Polle 2007	Sana 2015
Smoking										p=0.001*					
Open/conventional surgery					Difference in scores p=0.2	Difference in means p=0.51	Difference in median p=0.03*	Difference in median p=0.17						No significant differences	Multiple regression on (for laparoscopic approach) $\beta=0.331$ p=0.036*

Factor	Study												
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	1998	Dunker	2008	Eshuis 2010	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Perrin 2000
Increase d BMI													

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Factor	Study																		
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	Dunker 1998	2008	Eshuis	2010	Eshuis	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Ogden 2000	Perrin 2000	Polle 2007	Saha 2015	Scarpa 2009	Trindade 2017
Impaired QoL	r=0.52*	r=0.67*	r<0.41		r=0.5*							p<0.001*			r=0.51*		One unit increase in ASWAP score associated with a 0.62 decrease in IBDQ (p<0.0001).*		Psychological QoL r=0.56*
*Significant association found.																			

Factor	Study																	
	Abdovic	Bel 2015	Beld 2010	Chouliaras 2017	1998	Dunker	2008	Eshuis 2010	Kjaer 2014	Mauder 1999	McDermott 2015	Muller 2010	Ogden 2000	Perrin 2000	Polle 2007	Saha 2015	Scarpa 2009	Trindade 2017
<i>N.B: With some tools, higher scores indicate better body image/QoL and in others higher scores indicate worse body image/QoL. This may result in both positive and negative correlation coefficients. Where applicable, signs have been flipped for ease of interpretation to clearly show the positive correlation between body image and quality of life.</i>																		

Risk of Bias

The 31 studies relevant for questions 2-4 were assessed using criteria from the Joanna Briggs Institute critical appraisal tools for analytical cross-sectional and prevalence designs (Supplementary Table 2). Only cross-sectional data was relevant for the review. Poor reporting of quality criteria in many studies made quality assessment difficult. Where criteria were reported, the overall quality was variable. Most studies had some areas of low and higher quality. Only one study, McDermott (2015)⁽¹⁴⁾, was able to demonstrate adequate response rates, validated outcome measurement tools and adjustment for confounders. However, Chouliaras (2017)⁽³⁰⁾, Trindade (2017)⁽⁷²⁾, Lee (2015)⁽⁵¹⁾ and Bel (2015)⁽²³⁾ adjusted for confounders and used validated outcome measurement tools but lacked adequate response rates.

Twenty studies (64.5%) used an appropriate sample frame with acquisition of patients from outpatient settings, IBD registries or healthcare records. Eighteen studies (58.1%) clearly reported inclusion criteria applied when recruiting participants. Only twelve studies (38.7%) had response rates >75%. Fifteen studies (48.4%) used a tool which had been validated using factor analysis and internal consistency analysis to measure BI. The others used non-validated tools. Twelve studies^(14, 35, 50, 51, 59, 65, 66, 73) adjusted for potential confounders such as age, gender, BMI and previous surgery often using multiple regression models. Several studies reported limited demographic data. It should also be noted that sample sizes of many of the studies were small and confidence intervals were mostly not presented.

DISCUSSION

Summary of Findings

Overall, fifteen different tools were used across 57 studies to measure BI in IBD patients. These included QoL tools incorporating BI questions or domains, BI tools and other adapted questionnaires. None offer a defining threshold for presence or absence of BID, which is not commonly considered as a specific psychological disorder unlike body dysmorphia.

It remains unclear whether IBD patients suffer with BID more so than the general population as most studies reported mean values with no reference to healthy population values. Three studies estimated a prevalence of a negative BI based on one question and this varied between 21 and 81%. This wide variation likely reflects the differences in tools and study characteristics. All three studies were based on self-report questionnaires with a wide age range and registry or hospital-based population.

Certain factors including female gender, disease activity and steroid use were consistently found to be significantly associated with increased BID in IBD patients. There was also a significant association between increasing BID and decreasing QoL reported in eight studies. These findings are consistent with a previous narrative review⁽⁷⁹⁾ assessing BID and sexual functioning in IBD patients.

Strengths and Weaknesses of the Review

This is the first systematic review assessing BID in an IBD population, and a robust methodology was employed to ensure that bias and errors were minimised. A sensitive search strategy means that it is unlikely that relevant studies were missed and over 50 studies have contributed to the evidence base in an area previously unexplored by a systematic review.

The review has some limitations. Some of the extracted data is based on abstracts only where full texts could not be obtained from the authors. This will have resulted in some missing information.

Furthermore, qualitative studies were not included as this was considered beyond the scope of this review. It's likely that there are qualitative studies which could offer a deeper insight into perception of BI in IBD patients.

Strengths and Weaknesses of the Evidence

There are some weaknesses within the included evidence. All studies had some areas of high risk of bias or had poorly reported methodological criteria thus hampering quality assessment. Some studies had very low response rates leading to possible under-representation of certain groups. Few studies adjusted for confounders which could have resulted in overestimates of associations.

A further issue is the lack of healthy control groups. Although it appears that IBD patients are concerned about BI, it is difficult to determine whether they are

affected more than the general population. However, it has been found that children and adolescents with chronic illnesses such as asthma, cystic fibrosis and diabetes do have increased BID compared to healthy peers⁽⁸⁰⁾.

Non-validated tools were often used for measuring BI and the reliability and validity of findings based on these is therefore unknown. There is also still little known about potential changes in BI perception over time.

Findings in Context

This review is consistent with findings from the narrative review by Jedel (2015)⁽⁷⁹⁾ which found that BI could potentially be a problem in IBD patients. Whilst surgery has been found to be an important contributing factor in BID in other research⁽⁸¹⁾, it is unclear how it impacts upon IBD patients. An association between BID and poorer QoL has been highlighted in both.

Females and adolescents are more likely to be concerned with BI and to suffer with BID compared with males and older people⁽⁸²⁻⁸⁷⁾. Whilst we found inconsistent results surrounding age, IBD is often diagnosed in adolescence when BID could be more of a concern.

In oncology, BI is more widely researched. One study suggested gynaecologic cancer patients suffered with BID which predicted emotional well-being⁽⁸⁸⁾. Another study with advanced cancer patients suggested BID was associated with depression, anxiety and fatigue⁽⁸⁹⁾. Qualitative research in pregnancy⁽⁹⁰⁾ and systematic lupus⁽⁹¹⁾ suggests BI can affect medication compliance and that

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

445 patients would like more support around dealing with BI issues. This could also
446 be true for IBD patients.

447 Finally, a previous systematic review found that children with chronic conditions
448 were more likely to be dissatisfied with their body than healthy peers⁽⁹²⁾.
449 Although IBD patients were not included, patients with similar chronic diseases
450 like diabetes, cancer, asthma and scoliosis were, suggesting IBD patients could
451 be similarly affected.

452
453 **Implications**

454 This evidence identified in this review suggests an association between BID and
455 poorer QoL as well as finding factors influencing BI in IBD patients. There were
456 however limitations to the evidence in terms of methodological quality and/or
457 reporting. Also, results were difficult to compare across studies. More
458 promisingly, BI is becoming an increasingly assessed outcome, highlighting the
459 need for continued research in this area.

460 Current research suggests that age, gender, medication and disease activity in
461 IBD may impact upon BI. These could be taken into account by clinicians and
462 patients by altering therapy or targeting comorbidities which could have a
463 beneficial effect on BID. Interventions to improve BI could be incorporated into
464 treatment strategies, which may in turn help to improve QoL. A recent
465 systematic review⁽⁹³⁾ found that stress-management, mindfulness and talking
466 therapies may offer small to moderate improvements in BI however there is a
467 lack of evidence from good randomised controlled trials.

468 **Future Research**

469 Future research should focus on developing a consensus around which
470 validated tool or tools are best suited to measuring BID in an IBD population.
471 Whilst we describe validity of tools such as the Body Image Scale, we have not
472 independently verified this, therefore we could not recommend a particular tool.
473 Defining thresholds may allow estimation of the prevalence of BID in this
474 population. Establishing reference values in a healthy population would allow for
475 more meaningful interpretation of BID scores across different chronic diseases.
476 Enrolling patients from diagnosis and following them over time would be useful
477 to measure how BI changes with duration, activity of disease and treatment.
478 Whilst more severe IBD symptoms or invasive treatment options may
479 exacerbate BID, BID itself and any associated anxiety or depressive symptoms
480 may in turn exacerbate IBD symptoms^(94, 95), and future research should also
481 address this association. If BID is recognised and treated early it may contribute
482 to preventing worsening disease course. It may also be useful to encourage the
483 use of BI as a patient reported outcome in future IBD studies. This would
484 increase data on BID and lead to a greater understanding of the condition.

486 **CONCLUSION**

487 In conclusion, the evidence suggests a detrimental effect of IBD on BI, but
488 uncertainty remains due a lack of comparison data from healthy populations.
489 Associations of BID with disease-related factors such as steroid treatment,
490 fatigue, disease activity and surgery are apparent and findings suggest a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

correlation between impaired BI and poorer QoL. These results should be cautiously interpreted due to risk of bias and/or poor reporting of methodological criteria amongst included studies, and the wide variation between populations, BI tools, and scoring systems. Future studies should make use of validated measurement tools and include BI as a main outcome where appropriate.

Specific Author Contributions

SB identified the topic, undertook scoping, defined the question, developed the protocol and wrote the draft of the manuscript. IH contributed to the methods development and carried out second reviewer tasks as well as helping to draft, comment on and approve the final version of this paper. DM provided substantial methodological input to aid protocol development and assisted with drafting and reading, commenting on approving the final version. JD provided methodological input and read, commented on and edited the draft and approved the final version.

Financial Support

During this research Isobel Harris was funded by a National Institute for Health Research (NIHR) Research Methods - Systematic Review Fellowship and Sophie Beese was a locally funded trainee in systematic reviews at the University of Birmingham with agreement from the NIHR. This article presents independent research funded by the NIHR. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

513

514 **Acknowledgements**

515 Not applicable.

516 **Conflicts of Interest**

517 The authors have nothing to disclose.

REFERENCES

1. Centers for Disease Control and Prevention. What is inflammatory bowel disease (IBD)? 2014 [cited 2017 30/01/2017]. Available from: <https://www.cdc.gov/ibd/what-is-ibd.htm>.

2. Crohn's and Colitis UK. About Inflammatory Bowel Disease [cited 2017 18/02/2017]. Available from: <https://www.crohnsandcolitis.org.uk/about-inflammatory-bowel-disease>.

3. Kaplan GG. The global burden of IBD: from 2015 to 2025. *Nature Reviews Gastroenterology & Hepatology*. 2015;12:720-7.

4. National Eating Disorders Association. What is body image? [cited 2017 17/02/2017]. Available from: <https://www.nationaleatingdisorders.org/what-body-image>.

5. Griffiths S, Hay P, Mitchison D, Mond JM, McLean SA, Rodgers B, et al. Sex differences in the relationships between body dissatisfaction, quality of life and psychological distress. *Aust N Z J Public Health*. 2016;40:518-22.

6. Brown Z, Tiggemann M. Attractive celebrity and peer images on Instagram: Effect on women's mood and body image. *Body image*. 2016;19:37-43.

7. Cohen R, Blaszczynski A. Comparative effects of Facebook and conventional media on body image dissatisfaction. *Journal of eating disorders*. 2015;3:23.

8. Holland G, Tiggemann M. A systematic review of the impact of the use of social networking sites on body image and disordered eating outcomes. *Body image*. 2016;17:100-10.

9. Dyl J, Kittler J, Phillips KA, Hunt JI. Body Dysmorphic Disorder and Other Clinically Significant Body Image Concerns in Adolescent Psychiatric Inpatients: Prevalence and Clinical Characteristics. *Child Psychiatry and Human Development*. 2006;36:369-82.

10. Kim JS, Kang S. A Study on Body Image, Sexual Quality of Life, Depression, and Quality of Life in Middle-aged Adults. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2015;9:96-103.

11. Thomas F. Cash. Body Image Assessments [cited 2017 17/02/2017]. Available from: <http://www.body-images.com/assessments/>.

12. McDermott E, Moloney J, Rafter N, Keegan D, Byrne K, Doherty GA, et al. The body image scale: a simple and valid tool for assessing body image dissatisfaction in inflammatory bowel disease. *Inflamm Bowel Dis*. 2014;20:286-90.

13. NHS Choices. Corticosteroids - Side effects 2015 [updated 12/03/2015; cited 2017 19/02/2017]. Available from: [http://www.nhs.uk/Conditions/Corticosteroid-\(drugs\)/Pages/Sideeffects.aspx](http://www.nhs.uk/Conditions/Corticosteroid-(drugs)/Pages/Sideeffects.aspx).

14. McDermott E, Mullen G, Moloney J, Keegan D, Byrne K, Doherty GA, et al. Body image dissatisfaction: Clinical features, and psychosocial disability in inflammatory bowel disease. *Inflamm Bowel Dis*. 2015;21:353-60.

15. Zaghiyan K, Ghantiwala V, Le Q, Murrell Z, Fleshner P. Is body image and cosmesis better after double-port laparoscopic or open ileal pouch-anal anastomosis (IPAA)? *2011;54:e119*.

16. Michaela MB, Dianne N-S. Body dissatisfaction: an overlooked public health concern. *Journal of Public Mental Health*. 2014;13:64-9.

17. Bernstein CN, Hitchon CA, Walld R, Bolton JM, Sareen J, Walker JR, et al. Increased Burden of Psychiatric Disorders in Inflammatory Bowel Disease. *Inflamm Bowel Dis*. 2018.

18. NHS Choices. Inflammatory Bowel Disease 2015 [updated 20/03/2015; cited 2017 17/02/2017]. Available from: <http://www.nhs.uk/conditions/Inflammatory-bowel-disease/Pages/Introduction.aspx>.
19. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med*. 2009;6.
20. Beese S, Harris I, Dretzke J, Moore D. Body Image Dissatisfaction in Patients with Inflammatory Bowel Disease: A Systematic Review Protocol *BMC Systematic Reviews*. 2018.
21. The Joanna Briggs Institute. Joanna Briggs Institute Reviewers' Manual: 2016 edition. . 2016.
22. Abdovic S, Mocic Pavic A, Milosevic M, Persic M, Senecic-Cala I, Kolacek S. The IMPACT-III (HR) Questionnaire: A valid measure of health-related quality of life in Croatian children with inflammatory bowel disease. *Journal of Crohn's and Colitis*. 2013;7:908-15.
23. Bel LGJ, Vollebregt AM, Van der Meulen-de Jong AE, Fidder HH, Ten Hove WR, Vliet-Vlieland CW, et al. Sexual Dysfunctions in Men and Women with Inflammatory Bowel Disease: The Influence of IBD-Related Clinical Factors and Depression on Sexual Function. *J Sex Med*. 2015;12:1557-67.
24. Beld M, Van Balkom K, Visschers R, Van Gemert W, Breukink S. Long term results after restorative proctocolectomy with ileal pouch-anal anastomosis at young age. *Colorectal Dis*. 2010;12:16.
25. Bengtsson J, Lindholm E, Nordgren S, Berndtsson I, Oresland T, Borjesson L. Sexual function after failed ileal pouch-anal anastomosis. *Journal of Crohn's and Colitis*. 2011;5:407-14.
26. Brown C, Gibson PR, Hart A, Kaplan GG, Kachroo S, Ding Q, et al. Long-term outcomes of colectomy surgery among patients with ulcerative colitis. *Springerplus*. 2015;4:573.
27. Cabras PL, Giardinelli L, la Malfa GP, Galassi F. Variations in body image during autogenic training in patients with psychosomatic gastrointestinal disorders. *Giunti Organizzazioni Speciali*. 1986;178:27-33.
28. Camilleri-Brennan J, Munro A, Steele RJ. Does an ileoanal pouch offer a better quality of life than a permanent ileostomy for patients with ulcerative colitis? *J Gastrointest Surg*. 2003;7:814-9.
29. Carlsen K, Jakobsen C, Hansen LF, Paerregaard A, Kallemose T, Riis LB, et al. Quality of life in paediatric inflammatory bowel disease patients in a self-administered telemedicine randomised clinical study. *Journal of Crohn's and Colitis*. 2016;10:S421-S2.
30. Chouliaras G, Margoni D, Dimakou K, Fessatou S, Panayiotou I, Roma-Giannikou E. Disease impact on the quality of life of children with inflammatory bowel disease. *World J Gastroenterol*. 2017;23:1067-75.
31. de Rooy EC, Toner BB, Maunder RG, Greenberg GR, Baron D, Steinhart AH, et al. Concerns of patients with inflammatory bowel disease: results from a clinical population. *Am J Gastroenterol*. 2001;96:1816-21.
32. Drossman DA, Patrick DL, Mitchell CM, Zagami EA, Appelbaum MI. Health-related quality of life in inflammatory bowel disease. Functional status and patient worries and concerns. *Dig Dis Sci*. 1989;34:1379-86.
33. Dunker MS, Bemelman WA, Slors JF, van Duijvendijk P, Gouma DJ. Functional outcome, quality of life, body image, and cosmesis in patients after laparoscopic-assisted and conventional restorative proctocolectomy: a comparative study. *Dis Colon Rectum*. 2001;44:1800-7.
34. Dunker MS, Stiggelbout AM, van Hogezaand RA, Ringers J, Griffioen G, Bemelman WA. Cosmesis and body image after laparoscopic-assisted and open ileocolic resection for Crohn's disease. *Surg Endosc*. 1998;12:1334-40.

35. Eshuis EJ, Polle SW, Slors JF, Hommes DW, Sprangers MA, Gouma DJ, et al. Long-term surgical recurrence, morbidity, quality of life, and body image of laparoscopic-assisted vs. open ileocolic resection for Crohn's disease: a comparative study. *Dis Colon Rectum*. 2008;51:858-67.
36. Eshuis EJ, Slors JFM, Stokkers PCF, Sprangers MAG, Ubbink DT, Cuesta MA, et al. Long-term outcomes following laparoscopically assisted versus open ileocolic resection for Crohn's disease. *Br J Surg*. 2010;97:563-8.
37. Eshuis EJ, Voermans RP, Stokkers PCF, Van Berge Henegouwen MI, Fockens P, Bemelman WA. Laparoscopic resection with transcolonic specimen extraction for ileocaecal Crohn's disease. *Br J Surg*. 2010;97:569-74.
38. Gallo J, Grant A, Otley AR, Orsi M, Macintyre B, Gauvry S, et al. Do parents and children agree? Quality-of-life assessment of children with inflammatory bowel disease and their parents. *J Pediatr Gastroenterol Nutr*. 2014;58:481-5.
39. Giudici F, Scaringi S, Di Martino C, Ficari F, Bechi P. Rationalisation of the surgical technique for minimally invasive laparoscopic ileal pouch-anal anastomosis after previous total colectomy for ulcerative colitis. *J Minim Access Surg*. 2017;13:188-91.
40. Grant A, Kappelman M, Martin C, Otley A. A new domain structure for the impact-III, a pediatric inflammatory bowel disease (IBD) health reported quality of life (HRQOL) tool. *Inflamm Bowel Dis*. 2016;22:S7-S8.
41. Grant A, Otley A, Escher J, Hyams JS, Maa JF, Alperovich G, et al. Assessment of IMPACT III emotional and social functioning domain scores in adalimumab-treated paediatric patients with Crohn's disease. *Journal of Crohn's and Colitis*. 2016;10:S424-S5.
42. Grootenhuis MA, Maurice-Stam H, Derkx BH, Last BF. Evaluation of a psychoeducational intervention for adolescents with inflammatory bowel disease. *Eur J Gastroenterol Hepatol*. 2009;21:340-5.
43. Gudlaugsdottir K, Valsdottir EB, Stefansson TB. [Quality of Life after colectomy due to ulcerative colitis]. *Laeknabladid*. 2016;102:482-9.
44. Hagan M, Jambaulikar G, Osche-Gauvin K, Schwartz D, Higginbotham T, Cross R. Sexual function in patients with inflammatory bowel disease: Results of a web-based health survey. *Am J Gastroenterol*. 2014;109:S516.
45. Joachim G, Milne B. Inflammatory bowel disease: effects on lifestyle. *J Adv Nurs*. 1987;12:483-7.
46. Juan L, Ricardo DLV, Mayte V, Esther T. Gender differences in stoma-related quality of life in puerto ricans with IBD. *Am J Gastroenterol*. 2018;113 (Supplement 1):S14.
47. Kjaer MD, Laursen SB, Qvist N, Kjeldsen J, Poornoroozy PH. Sexual function and body image are similar after laparoscopy-assisted and open ileal pouch-anal anastomosis. *World J Surg*. 2014;38:2460-5.
48. Knowles SR, Gass C, Macrae F. Illness perceptions in IBD influence psychological status, sexual health and satisfaction, body image and relational functioning: A preliminary exploration using Structural Equation Modeling. *Journal of Crohn's and Colitis*. 2013;7:e344-e50.
49. Knowles SR, Wilson J, Wilkinson A, Connell W, Salzberg M, Castle D, et al. Psychological well-being and quality of life in Crohn's disease patients with an ostomy: a preliminary investigation. *Journal of Wound, Ostomy, & Continence Nursing*. 2013;40:623-9.
50. Kuruvilla K, Osler T, Hyman NH. A comparison of the quality of life of ulcerative colitis patients after IPAA vs ileostomy. *Dis Colon Rectum*. 2012;55:1131-7.
51. Lee D, Baldassano RN, Otley AR, Albenberg L, Griffiths AM, Compher C, et al. Comparative effectiveness of nutritional and biological therapy in North American children with active Crohn's disease. *Inflamm Bowel Dis*. 2015;21:1786-93.

52. Liwanag MJ, Liu JX, Tan LN, Huang JG, Quak SH, Aw MM. Health related quality of life in paediatric inflammatory bowel disease in a Southeast Asian population. *Journal of Crohn's and Colitis*. 2014;8:S409.
53. Mason A, Malik S, McMillan M, McNeilly JD, Bishop J, McGrogan P, et al. A prospective longitudinal study of growth and pubertal progress in adolescents with inflammatory bowel disease. *Horm Res Paediatr*. 2015;83:45-54.
54. Maunder R, Toner B, De Rooy E, Moskovitz D. Influence of sex and disease on illness-related concerns in inflammatory bowel disease. *Can J Gastroenterol*. 1999;13:728-32.
55. Mazzoccone A, et al. A study of body image in patients with chronic colon and liver diseases. *Giunti Organizzazioni Speciali*. 1980;155-156:105-13.
56. McDermott E, Moloney J, Rafter N, Keegan D, Byrne K, Doherty GA, et al. The body image scale: A simple and valid tool for assessing body image dissatisfaction in inflammatory bowel disease. *Inflamm Bowel Dis*. 2014;20:286-90.
57. Mountifield R, Bampton P, Prosser R, Muller K, Andrews JM. Fear and fertility in inflammatory bowel disease: A mismatch of perception and reality affects family planning decisions. *Inflamm Bowel Dis*. 2009;15:720-5.
58. Mukhopadhyay A, Probert S, Smith C, Maville C, Renji E, Bunn S, et al. IMPACT III- disease-specific health-related quality of life (HRQOL) for children with Crohn's disease (CD) on infliximab-a single centre experience. *J Pediatr Gastroenterol Nutr*. 2017;64 (Supplement 1):519-20.
59. Muller KR, Prosser R, Bampton P, Mountifield R, Andrews JM. Female gender and surgery impair relationships, body image, and sexuality in inflammatory bowel disease: Patient perceptions. *Inflamm Bowel Dis*. 2010;16:657-63.
60. Navas-Lopez VM, Martin-De-Carpi J, Grant A, Walters TD, Ruemmele F, Mack D, et al. Quality of life in paediatric Crohn's disease: Data from the Imagekids study. *Journal of Crohn's and Colitis*. 2016;10:S145-S6.
61. Ogden CA, Akobeng AK, Abbott J, Aggett P, Sood MR, Thomas AG. Validation of an instrument to measure quality of life in British children with inflammatory bowel disease. *J Pediatr Gastroenterol Nutr*. 2011;53:280-6.
62. Perrin JM, Kuhlthau K, Chughtai A, Romm D, Kirschner BS, Ferry GD, et al. Measuring quality of life in pediatric patients with inflammatory bowel disease: Psychometric and clinical characteristics. *J Pediatr Gastroenterol Nutr*. 2008;46:164-71.
63. Plevinsky JM, Greenley RN. Exploring health-related quality of life and social functioning in adolescents with inflammatory bowel diseases after attending camp oasis and participating in a facebook group. *Inflamm Bowel Dis*. 2014;20:1611-7.
64. Polle SW, Dunker MS, Slors JF, Sprangers MA, Cuesta MA, Gouma DJ, et al. Body image, cosmesis, quality of life, and functional outcome of hand-assisted laparoscopic versus open restorative proctocolectomy: long-term results of a randomized trial. *Surg Endosc*. 2007;21:1301-7.
65. Ponsioen CY, de Groof EJ, Eshuis EJ, Gardenbroek TJ, Bossuyt PMM, Hart A, et al. Laparoscopic ileocaecal resection versus infliximab for terminal ileitis in Crohn's disease: a randomised controlled, open-label, multicentre trial. *The Lancet Gastroenterology & Hepatology*. 2017;2:785-92.
66. Saha S, Zhao YQ, Shah SA, Esposti SD, Lidofsky S, Bright R, et al. Body image dissatisfaction in patients with inflammatory bowel disease. *Inflamm Bowel Dis*. 2015;21:345-52.
67. Savarino JR, Venkatesh RD, Israel EJ, Kaplan JL. Health-related quality of life in pediatric inflammatory bowel disease patients receiving infliximab: A pilot study using the impact-III questionnaire. *J Pediatr Gastroenterol Nutr*. 2016;63:S362-S3.

68. Scarpa M, Ruffolo C, Bassi D, Boetto R, D'Inca R, Buda A, et al. Intestinal surgery for Crohn's disease: Predictors of recovery, quality of life, and costs. *J Gastrointest Surg*. 2009;13:2128-35.
69. Shah S, Urban M, Gracely E, Nandi N. Anonymous self perception survey of sexuality and body image in inflammatory bowel disease. *Am J Gastroenterol*. 2017;112 (Supplement 1):S377-S9.
70. Shepanski MA, Hurd LB, Culton K, Markowitz JE, Mamula P, Baldassano RN. Health-related quality of life improves in children and adolescents with inflammatory bowel disease after attending a camp sponsored by the Crohn's and Colitis Foundation of America. *Inflamm Bowel Dis*. 2005;11:164-70.
71. Swedish E, Blucker RT, Grunow J, Suorsa K, Jacobs NJ. Severity of illness and quality of life over time in pediatric inflammatory disease patients. *Gastroenterology*. 2015;1):S635.
72. Trindade IA, Ferreira C, Pinto-Gouveia J. The effects of body image impairment on the quality of life of non-operated Portuguese female IBD patients. *Qual Life Res*. 2017;26:429-36.
73. Vlahou CH, Cohen LL, Woods AM, Lewis JD, Gold BD. Age and body satisfaction predict diet adherence in adolescents with inflammatory bowel disease. *J Clin Psychol Med Settings*. 2008;15:278-86.
74. Weinryb RM, Gustavsson JP, Barber JP. Personality predictors of dimensions of psychosocial adjustment after surgery. *Psychosom Med*. 1997;59:626-31.
75. Weinryb RM, Gustavsson JP, Barber JP. Personality traits predicting long-term adjustment after surgery for ulcerative colitis. *J Clin Psychol*. 2003;59:1015-29.
76. Werner H, Landolt MA, Buehr P, Koller R, Nydegger A, Spalinger J, et al. Validation of the IMPACT-III quality of life questionnaire in Swiss children with inflammatory bowel disease. *Journal of Crohn's and Colitis*. 2014;8:641-8.
77. Zambonin D, Giudici F, Ficari F, Rogai F, Scaringi S. Short-and long-term outcome of minimally invasive approach for Crohn's disease: Comparison between single incision, roboticassisted and conventional laparoscopy. *Journal of Crohn's and Colitis*. 2018;12 (Supplement 1):S411-S2.
78. Kjaer MD, Laursen SB, Poornorozy PH. Sexual function and body image is similar after laparoscopic and open ileal pouch-anal anastomosis. *Gastroenterology*. 2014;1):S-383.
79. Jedel S, Hood MM, Keshavarzian A. Getting personal: a review of sexual functioning, body image, and their impact on quality of life in patients with inflammatory bowel disease. *Inflamm Bowel Dis*. 2015;21:923-38.
80. Pinquart M. Body image of children and adolescents with chronic illness: A meta-analytic comparison with healthy peers. *Body Image*. 2013;10:141-8.
81. Allison M, Lindsay J, Gould D, Kelly D. Surgery in young adults with inflammatory bowel disease: a narrative account. *International journal of nursing studies*. 2013;50:1566-75.
82. Rosenblum GD, Lewis M. The relations among body image, physical attractiveness, and body mass in adolescence. *Child development*. 1999;70:50-64.
83. Bearman SK, Martinez E, Stice E, Presnell K. The Skinny on Body Dissatisfaction: A Longitudinal Study of Adolescent Girls and Boys. *Journal of youth and adolescence*. 2006;35:217-29.
84. Brennan M, Lalonde C, Bain J. Body Image Perceptions: Do Gender Differences Exist? *Psi Chi J Undergrad Res*. 2010.
85. Miranda VP, Conti MA, de Carvalho PH, Bastos RR, Ferreira ME. Body image in different periods of adolescence. *Rev Paul Pediatr*. 2014;32:63-9.
86. Ando S, Osada H. Age and gender differences in body image over the life span: Relationships between physical appearance, health and functioning 2009. 1-16 p.
87. Prais TA, Janowsky JS. Assessment of body image in younger and older women. *J Gen Psychol*. 2010;137:225-38.


- 1
2
3
4
5 759 88. Teo I, Cheung YB, Lim TYK, Namuduri RP, Long V, Tewani K. The relationship between
6 760 symptom prevalence, body image, and quality of life in Asian gynecologic cancer patients.
7 761 *Psycho-oncology*. 2017.
8 762 89. Rhondali W, Chisholm GB, Daneshmand M, Allo J, Kang DH, Filbet M, et al. Association
9 763 between body image dissatisfaction and weight loss among patients with advanced cancer and
10 764 their caregivers: a preliminary report. *Journal of pain and symptom management*.
11 765 2013;45:1039-49.
12 766 90. Watson B, Broadbent J, Skouteris H, Fuller-Tyszkiewicz M. A qualitative exploration of
13 767 body image experiences of women progressing through pregnancy. *Women Birth*. 2016;29:72-
14 768 9.
15 769 91. Hale ED, Radvanski DC, Hassett AL. The man-in-the-moon face: a qualitative study of
16 770 body image, self-image and medication use in systemic lupus erythematosus. *Rheumatology*.
17 771 2015;54:1220-5.
18 772 92. Piquart M. Body image of children and adolescents with chronic illness: a meta-
19 773 analytic comparison with healthy peers. *Body Image*. 2013;10:141-8.
20 774 93. Alleva JM, Sheeran P, Webb TL, Martijn C, Miles E. A Meta-Analytic Review of Stand-
21 775 Alone Interventions to Improve Body Image. *PLoS One*. 2015;10:e0139177.
22 776 94. Mikocka-Walus A, Pittet V, Rossel J-B, von Känel R, Anderegg C, Bauerfeind P, et al.
23 777 Symptoms of Depression and Anxiety Are Independently Associated With Clinical Recurrence
24 778 of Inflammatory Bowel Disease. *Clin Gastroenterol Hepatol*. 2016;14:829-35.e1.
25 779 95. Kochar B, Barnes EL, Long MD, Cushing KC, Galanko J, Martin CF, et al. Depression Is
26 780 Associated With More Aggressive Inflammatory Bowel Disease. *The American Journal Of*
27 781 *Gastroenterology*. 2017;113:80.
28
29
30
31 782
32
33
34 783
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

784 **Figures**

785 **1.** The selection process of records for inclusion/exclusion detailed in a PRISMA
786 flowchart.

Confidential: For Review Only



437x618mm (96 x 96 DPI)

Supplementary Data Content - Table 1

MEDLINE Search Strategy – OVID MEDLINE In process & other non-indexed citations and OVID MEDLINE.

Search	Query
#1	exp inflammatory bowel diseases/
#2	inflammatory bowel disease*.mp.
#3	exp Colitis, Ulcerative/
#4	ulcerative colitis.mp.
#5	exp Crohn disease/
#6	Crohn* disease.mp.
#7	Crohn*.mp.
#8	IBD.mp.
#9	CD.mp.
#10	UC.mp.
#11	1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10
#12	exp body image/
#13	body image.mp.
#14	body dissatisfaction.mp.
#15	body awareness.mp.
#16	body concern*.mp.
#17	body attitude*.mp.
#18	body preoccupation.mp.

#19	body perception.mp.
#20	body anxiety.mp.
#21	body conscious*.mp.
#22	12 OR 13 OR 14 OR 15 OR 16 OR 17 OR 18 OR 19 OR 20 OR 21
#23	11 AND 22

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

1 **Supplementary Data Content - Table 2** Results of risk of bias assessment using Joanna Briggs Institute Tools

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Abdovic et al (2013)	Yes	Yes	Yes	Yes	N/A	No	No	Yes	Yes	Yes
Bel et al (2015)	Yes	Yes	Yes	Yes	N/A	Yes	Yes	Yes	Yes	No
Beld et al (2010)	Yes	Yes	No	Yes	Yes	No	No	No	Yes	Yes

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Bengtsson et al (2011)	Yes	No	No	Yes	Yes	Unclear	No	Yes	Yes	No
Brown et al (2015)	Yes	Yes	Yes	Yes	Yes	Unclear	No	No	Yes	No
Chouliaras et al (2017)	Yes	Yes	Yes	Yes	N/A	Yes	Yes	Yes	Yes	Unclear
de Rooy et al (2001)	Unclear	No	Yes	No	N/A	Yes	Yes	Yes	Yes	Unclear

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Dunker et al (1998)	No	No	Yes	Unclear	Unclear	No	No	No	Yes	Yes
Dunker et al (2001)	Unclear	No	Unclear	Yes	Unclear	Unclear	Unclear	No	Yes	Yes
Eshuis et al (2008)	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes
Eshuis et al	Unclear	Yes	Unclear	Yes	Yes	No	No	No	Yes	Yes

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
(2010)										
Gallo et al (2014)	Yes	Yes	Yes	Yes	N/A	No	No	Yes	Yes	Yes
(2014)										
Giudici et al (2017)	No	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
(2017)										
Grootenhuus (2009)	Yes	Yes	Yes	N/A	Unclear	Yes	Yes	No	Yes	No
(2009)										

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Kjaer et al (2014)	Unclear	Unclear	Yes	Yes	Unclear	No	No	No	Yes	No
Kuruvilla et al (2012)	Yes	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear	Unclear
Lee et al (2015)	Yes	Yes	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	Unclear
Mason et al (2015)	Yes	Yes	Yes	Yes	N/A	No	No	Yes	Yes	Unclear

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Maunder et al (1999)	Unclear	Unclear	Yes	Yes	N/A	No	No	Yes	Yes	Unclear
McDermott et al (2015)	Yes	Yes	Unclear	Unclear	N/A	Yes	Yes	Yes	Yes	Yes
Muller et al (2010)	Yes	Yes	Unclear	Unclear	N/A	Yes	Yes	No	Yes	Unclear
Ogden et al	Unclear	Unclear	Unclear	Unclear	N/A	N/A	N/A	Yes	Yes	Yes

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
(2011)										
Perrin et al (2008)	Yes	Yes	Yes	Unclear	N/A	N/A	N/A	Yes	Yes	No
Polle et al (2007)	Unclear	Unclear	Unclear	Yes	Unclear	Yes	No	No	Yes	Yes
Ponsioen et al (2017)	Yes	Yes	Yes	Yes	Unclear	Unclear	Unclear	Yes	Unclear	No

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Saha et al (2015)	Yes	Yes	Yes	Yes	N/A	Yes	Yes	Unclear	Yes	No
Scarpa et al (2009)	Unclear	Yes	Unclear	Yes	Yes	Yes	Yes	No	Yes	Yes
Shepanksi (2005)	Yes	Unclear	Unclear	Yes	Unclear	Unclear	No	Yes	Yes	No
Trindade et al (2017)	Yes	Yes	Yes	Unclear	N/A	Yes	Yes	Yes	Yes	Unclear

Study	Was the sample frame appropriate to address the target population?	Were the criteria for inclusion in the sample clearly defined?	Were the study subjects and the setting described in detail?	Was the exposure measured in a valid and reliable way?	Were objective, standard criteria used for measurement of the condition?	Were confounding factors identified?	Were strategies to deal with confounding factors stated?	Were the outcomes measured in a valid and reliable way?	Was appropriate statistical analysis used?	Was the response rate adequate, and if not, was the low response rate managed appropriately? (>75%)
Vlahou et al (2008)	Yes	Unclear	No	Unclear	N/A	Yes	Yes	No	Yes	Unclear
Voermans et al (2010)	Unclear	Yes	Unclear	Yes	N/A	No	No	No	Yes	Yes

Confidential: For Review Only